

Localization of Neurological Lesions

Chong-hao Zhao, MD, PhD

*Board Certified in Neurology, Headache Medicine,
Pain Medicine, Medical Acupuncture*

State of California X-ray Fluoroscopy Supervisor and Operator

Voluntary Teaching Faculty for Graduate Orofacial Pain Program at UCLA


Property of Chonghao Zhao,



California Headache and Pain Center
420 W. Las Tunas Dr., San Gabriel, CA 91776
Tel: 626-457-1688; www.chpci.com
Email: Drzhao@chpci.com

Brain Lesions Suggested by Abnormal Mental Status and Speech

- Mental status: cortex dysfunction - confusion, lethargy/coma
- Speech:
 - Aphasia:
 - Broca's (motor) aphasia: preserved comprehension, non-fluent speech. Lesion to the dominant (left) hemisphere: **inferior frontal gyrus** of the frontal lobe.
 - Wernicke's aphasia (sensory): poor comprehension, fluent but often meaningless speech. Lesion to the **supramarginal gyrus** of the parietal lobe and upper part of temporal lobe.





Other Speech Problems

- Speech:
 - Dysphonia: unable to produce normal volume of sound or speaks in a whisper. Lesion of vocal cord, laryngeal problem, myasthenia gravis
 - Dysarthria: slurred speech, lesion in cerebellum, upper and lower motor neuron disease, parkinsonism



Brain Lesions Suggested by the Abnormal Signs

- Meningeal irritation:
 - Neck stiffness
 - Brudzinski sign: lift the head and look for hip and knee flexion.
 - Positive: hip and knee flexion
 - Kernig's sign: flex the leg at the hip with the knee flexed, and try to extend the knee.
 - Positive: resistance to knee straightening




Lesions Suggested by Abnormal Cranial Nerve Function

- CN 1: loss of smelling
- CN 2: loss of visual acuity
- CN 3,4,6: loss of extraocular eye movement
- CN 5: abnormal facial sensation, loss of corneal reflex and symmetrical jaw movement
- CN 7: ipsilateral facial weakness
- CN 8: hearing loss
- CN 9, 10: loss of palate elevation and gag reflex
- CN 11: loss of shoulder shrugging function
- CN 12: tongue deviation to the affected side






Lesions of Motor, Coordination, and Sensory Pathways (Overview)




Lesions of Motor Pathway – The Descending Corticospinal Tract

- Lesion above the medulla pyramidal decussation:
 - E.g. motor cortex (frontal lobe), internal capsule, midbrain, pon, medulla, etc
 - Muscle weakness contralateral to the side of lesion
 - Babinski's sign: positive, contralateral to the side of lesion
- Lesion below the decussation:
 - E.g. cervical cord
 - Muscle weakness ipsilateral to the side of lesion
 - Babinski's sign: positive, ipsilateral to the side of lesion





Lesions of the Coordination Pathway – The Spinocerebellar Tract

- Transmitted from the spinal cord to the same side of the cerebellum
- Ataxia to the same side of the lesion in the coordination test.




Lesions of the Sensory Pathway – The Spinothalamic Tract



- Loss of pain and temperature on the contralateral side, beginning 1-2 level below to the lesion (decussation at the anterior white commissure usually occurs 1-2 spinal nerve segments above the point of entry)
- Lateral STT (pain and temperature);
Anterior STT (pressure and crude touch)




Lesions of Sensory Pathway – The Dorsal Column - Medial Lemniscus Pathway



Dorsal Surface of Medulla Oblongata

- **Nucleus gracilis & nucleus cuneatus** = sensory neurons
- relay information to thalamus on opposite side of brain
- **5 cranial nerves** arise from medulla -- 8 thru 12




- Dorsal column (in spinal cord, synapse in Nu. cuneatus and gracilis before decussation) - medial lemniscus (in brainstem) tract
 - **Below** the sensory decussation in the medulla (lesion of the dorsal column): loss of light touch, 2-point discrimination, vibration, and proprioception/joint positional sense to the **ipsilateral** side, and below the level of the lesion
 - **Above** the decussation (lesion of the medial lemniscus): sensory loss **contralateral** to the lesion



Lesions Suggested by the Abnormal Gait

- Stroke or lesion to the frontal motor cortex: hemiplegic gait contralateral to the lesion
- Cerebellar ataxia: deviate to the ipsilateral side of the lesion
- Loss of posterior column function: sensory ataxia, loss of proprioception/joint positional sense in Romberg's test
 - Above the sensory decussation in Medulla: contralateral to lesion
 - Below: ipsilateral
- Basal ganglion
 - components: dorsal striatum (caudate nucleus & putamen), ventral striatum (nucleus accumbens and olfactory tubercle), globus pallidus, ventral pallidum, substantia nigra, subthalamic nucleus
 - dysfunction: shuffle gait in Parkinson's disease.



Suggested Readings:

- Memorix Neurology by Peter Berlitz, 1996
- Neurological Examination Made Easy by Geraint Fuller, 1995
- Clinical Neurological Neuroanatomy Made Ridiculously Simple, by Stephen Goldberg, 1990

